

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
3 April 2003 (03.04.2003)

PCT

(10) International Publication Number  
WO 03/027239 A2

(51) International Patent Classification<sup>7</sup>: C12N

(21) International Application Number: PCT/US02/29931

(22) International Filing Date:  
20 September 2002 (20.09.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/325,129 26 September 2001 (26.09.2001) US

(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HOWARD, Andrew, D. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). PAN, Jie [CN/US]; 126 East Lincoln

Avenue, Rahway, NJ 07065-0907 (US). FONG, Tung, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). MARSH, Donald, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). SAILER, Andreas, W. [DE/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(81) Designated States (national): CA, JP, US.


(84) Designated States (regional): European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MELANIN-CONCENTRATING HORMONE RECEPTOR ANTAGONIST BINDING PROTEIN



(57) Abstract: The present invention features MCH-1R antagonist binding proteins. MCH-1R antagonist binding proteins described herein are based on an MCH-1R having one or more alterations to the second intracellular loop or carboxy terminus that render the receptor substantially inactive to MCH binding. An MCH-1R antagonist binding protein can bind MCH-1R antagonists, but does not exhibit high affinity MCH binding and is not activated by the MCH.

## TITLE OF THE INVENTION

MELANIN-CONCENTRATING HORMONE RECEPTOR ANTAGONIST  
BINDING PROTEIN

## 5 CROSS-REFERENCE TO RELATED APPLICATIONS

The present application claims priority to provisional application U.S. Serial No. 60/325,129, filed September 26, 2001, hereby incorporated by reference herein.

## 10 BACKGROUND OF THE INVENTION

The references cited in the present application are not admitted to be prior art to the claimed invention.

Neuropeptides present in the hypothalamus play a major role in mediating the control of body weight. (Flier, *et al.*, 1998. *Cell*, 92, 437-440.)

- 15 Melanin-concentrating hormone (MCH) is a cyclic 19-amino acid neuropeptide synthesized as part of a larger pre-prohormone precursor in the hypothalamus which also encodes neuropeptides NEI and NGE. (Nahon, *et al.*, 1990. *Mol. Endocrinol.* 4, 632-637.) MCH was first identified in salmon pituitary, and in fish MCH affects melanin aggregation thus affecting skin pigmentation. In trout and in eels MCH has  
20 also been shown to be involved in stress induced or CRF-stimulated ACTH release. (Kawauchi, *et al.*, 1983. *Nature* 305, 321-323.)

- In humans two genes encoding MCH have been identified that are expressed in the brain. (Breton, *et al.*, 1993. *Mol. Brain Res.* 18, 297-310.) In mammals MCH has been localized primarily to neuronal cell bodies of the  
25 hypothalamus which are implicated in the control of food intake, including perikarya of the lateral hypothalamus and zona inertia. (Knigge, *et al.*, 1996. *Peptides* 17, 1063-1073.)

- Pharmacological and genetic evidence suggest that the primary mode of MCH action is to promote feeding (orexigenic). MCH mRNA is up regulated in  
30 fasted mice and rats and in the *ob/ob* mouse. (Qu, *et al.*, 1996. *Nature* 380, 243-247.) Injection of MCH centrally (ICV) stimulates food intake and MCH antagonizes the hypophagic effects seen with  $\alpha$  melanocyte stimulating hormone ( $\alpha$ MSH). (Qu, *et al.*, 1996. *Nature* 380, 243-247.) MCH deficient mice are lean, hypophagic and have increased metabolic rate. (Shimada, *et al.*, 1998. *Nature* 396, 670-673.) Transgenic

mice overexpressing MCH are hyperphagic and develop insulin resistance and mild obesity. (Ludwig, *et al.*, 2001, *J. Clin. Invest.* 107, 379-386.)

MCH action is not limited to modulation of food intake as effects on the hypothalamic-pituitary-axis have been reported. (Nahon, 1994. *Critical Rev. in Neurobiol.* 8, 221-262.) MCH can modulate stress-induced release of ACTH. (Nahon, 1994. *Critical Rev. in Neurobiol.* 8, 221-262.)

Several references describe a human melanin-concentrating hormone receptor ("MCH-1R"). (Chambers, *et al.*, 1999. *Nature* 400, 261-265, Saito, *et al.*, 1999. *Nature* 400, 265-269, Bächner, *et al.*, 1999. *FEBS Letters* 457:522-524, Shimomura, *et al.*, 1999. *Biochemical and Biophysical Research Communications* 261, 622-626.)

#### SUMMARY OF THE INVENTION

The present invention features MCH-1R antagonist binding proteins. MCH-1R antagonist binding proteins described herein are based on an MCH-1R having one or more alterations to the second intracellular loop or carboxy terminus that render the receptor substantially inactive to MCH binding. An MCH-1R antagonist binding protein can bind MCH-1R antagonists, but does not exhibit high affinity MCH binding and is not activated by the MCH.

Thus, a first aspect of the present invention describes an MCH-1R antagonist binding protein selected from the group consisting of:

- a) a MCH-1R antagonist binding protein having one or more alterations in the second intracellular loop region that render MCH-1R substantially inactive to MCH binding; and
- b) a MCH-1R antagonist binding protein having one or more alterations in the C-terminal that render MCH-1R substantially inactive to MCH binding.

"Substantially inactive to MCH binding" indicates that MCH binding, if present, is up to about 10% the level of binding to human MCH-1R. In different embodiments binding is 5% or less, and undetectable.

Another aspect of the present invention describes a nucleic acid comprising a nucleotide sequence encoding an MCH-1R antagonist binding protein. In an embodiment of the present invention, the nucleic acid is an expression vector.

Another aspect of the present invention describes a recombinant cell comprising an expression vector encoding an MCH-1R antagonist binding protein.

The nucleotide sequence encoding the MCH-1R antagonist binding protein is functionally coupled to a promoter recognized by the cell.

Another aspect of the present invention describes a method of screening for a compound able to bind an MCH-1R antagonist binding protein. The method involves contacting an MCH-1R antagonist binding protein with the compound and measuring the ability of the compound to bind to the protein.

Another aspect of the present invention describes a method of preparing a MCH-1R antagonist binding protein. The method involves growing a recombinant cell containing an expression vector encoding an MCH-1R antagonist binding protein.

Other features and advantages of the present invention are apparent from the additional descriptions provided herein including the different examples. The provided examples illustrate different components and methodology useful in practicing the present invention. The examples do not limit the claimed invention. Based on the present disclosure the skilled artisan can identify and employ other components and methodology useful for practicing the present invention.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates [ $^{125}$ I]Phe $^{13}$ Tyr $^{19}$ -MCH binding to MCH-1R(R141H) and MCH-1R(R155A).

Figure 2 illustrates [ $^{125}$ I]Phe $^{13}$ Tyr $^{19}$ -MCH binding to MCH-1R( $\Delta$ 316/EGFP).

Figure 3 illustrates the lack of functional activation of MCH-1R(R141H) by MCH. Functional activation was assayed by measuring mobilization of intracellular calcium.

Figure 4 illustrates the lack of functional activation of MCH-1R(R155A) by MCH. Functional activation was assayed by measuring mobilization of intracellular calcium.

Figure 5 illustrates the lack of functional activation of MCH-1R(i2MC4R) by MCH. Functional activation was assayed by measuring mobilization of intracellular calcium.

Figure 6 illustrates the lack of functional activation of MCH-1R( $\Delta$ 316/EGFP) by MCH. Functional activation was assayed by measuring mobilization of intracellular calcium.

## DETAILED DESCRIPTION OF THE INVENTION

Directed mutagenesis of the human MCH-1R has generated MCH-1R antagonist binding proteins that selectively bind MCH-1R antagonists. The MCH-1R antagonists do not exhibit high affinity MCH agonist binding and are not activated by MCH. Uses of MCH-1R antagonist binding proteins include screening for potential receptor antagonists and studying protein trafficking.

Different types of MCH-1R antagonist binding proteins were obtained by altering MCH-1R in the second intracellular loop region and by deleting a portion of the carboxy terminus. Alterations to the second intracellular loop region to produce a MCH-1R antagonist binding protein include single and multiple amino acid changes.

The MCH-1R antagonist binding proteins MCH-1R(R141H) and MCH-1R(R155A) contain single amino acid changes in the second intracellular loop region of MCH-1R. The amino acid sequences of MCH-1R(R141H) and MCH-1R(R155A) are provided by SEQ. ID. NO. 1 and SEQ. ID. NO. 2.

Position 141 is within the highly conserved DRY signature sequence found in most G-protein coupled receptors. The DRY signature sequence has been suggested to be involved in G-protein interaction. (Rosenthal, *et al.*, *J. Biol. Chem.* 268:13030-3, 1993.)

The MCH-1R antagonist binding protein MCH-1R(i2/MC4R) contains the MCH-1R, except the second intracellular loop which is replaced by the corresponding second intracellular loop of human MC4R. The amino acid sequence of MCH-1R(i2/MC4R) is provided by SEQ. ID. NO. 3.

MC4R is the melanocortin-4 receptor. (Yang *et al.*, *Biochemistry* 39: 14900-11, 2000, Gantz *et al.*, *J. Biol. Chem.* 268:15174-9, 1993.) Alterations to MC4R are described, for example, by Fraendberg, *et al. Biochem. Biophys. Res. Commun.* 245:490-492, 1998.

An example of a C-terminal deletion is provided by MCH-1R( $\Delta$ 316/EGFP) where the C-terminal 37 amino acids of MCH-1R was deleted and the enhanced green fluorescence protein (EGFP) was added to the C-terminus. C-terminal deletions to the human somatostatin receptor type 5 have been described by Hukovic, *et al. Journal of Biological Chemistry* 273:21416-21422, 1998.

The amino acid sequence of MCH-1R( $\Delta$ 316/EGFP) is provided by SEQ. ID. NO. 4. The EGFP sequence facilitates the study of protein trafficking.

### Production of MCH-1R Antagonist Binding Protein

Different MCH-1R antagonist binding protein can be obtained based on the guidance provided herein. The provided guidance includes the identification of particular mutations and regions useful for producing MCH-1R binding antagonists.

- 5 Preferred MCH-1R antagonist binding proteins are based on the human MCH-1R sequence.

MCH-1R antagonist binding protein should be able to bind an MCH antagonist, but not MCH. Different MCH-1R antagonist binding proteins can be produced, for example, by starting with an MCH-1R antagonist binding protein described herein and making additional alterations.

Alterations to a polypeptide not expected to alter polypeptide functioning can be made taking into account amino acid R groups. Differences in naturally occurring amino acids are due to different R groups. An R group affects different properties of an amino acid such as physical size, charge, and hydrophobicity. Amino acids can be divided into different groups as follows: neutral and hydrophobic (alanine, valine, leucine, isoleucine, proline, tryptophan, phenylalanine, and methionine); neutral and polar (glycine, serine, threonine, tyrosine, cysteine, asparagine, and glutamine); basic (lysine, arginine, and histidine); and acidic (aspartic acid and glutamic acid).

20 Generally, in substituting different amino acids to maintain antagonist binding it is preferable to exchange amino acids having similar properties. Substituting different amino acids within a particular group, such as substituting valine for leucine, arginine for lysine, and asparagine for glutamine are good candidates for not causing a change in antagonist binding.

25 In different embodiments MCH-1R antagonist binding protein: (1) have a sequence similarity of at least about 90%, preferably at least about 95% with either SEQ. ID. NOs. 1, 2, 3, 4, or a human MCH-1R with a deletion of about 37 amino acids; or (2) provide a sequence with up to about 20 alterations from SEQ. ID. NOs. 1, 2, 3, 4, or a human MCH-1R with a deletion of about 37 amino acids.

30 Sequence similarity for polypeptides can be determined by the BLAST. (Altschul, *et al.*, 1997. *Nucleic Acids Res.* 25, 3389-3402, hereby incorporated by reference herein.) In one embodiment sequence similarity is determined using tBLASTn search program with the following parameters: MATRIX:BLOSUM62, PER RESIDUE GAP COST: 11, and Lambda ratio: 1.

Alterations to amino acid sequences are additions, deletions, and substitutions. In different embodiments the MCH-1R polypeptide has 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or 10-20, alterations from SEQ. ID. NOs. 1, 2, 3, 4, or a human MCH-1R with a deletion of about 37 amino acids.

5 MCH-1R antagonist binding protein can be synthesized using standard techniques including those involving chemical synthesis and those involving biochemical synthesis. Techniques for chemical synthesis of polypeptides are well known in the art. (See *e.g.*, Vincent, in *Peptide and Protein Drug Delivery*, New York, N.Y., Dekker, 1990.)

10 Biochemical synthesis techniques for polypeptides are also well known in the art. Such techniques employ a nucleic acid template for polypeptide synthesis. Examples of techniques for introducing nucleic acid into a cell and expressing the nucleic acid to produce protein are provided in references such as Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 1987-1998, and Sambrook, *et al.*,  
15 *Molecular Cloning, A Laboratory Manual*, 2<sup>nd</sup> Edition, Cold Spring Harbor Laboratory Press, 1989.

Starting with a particular amino acid sequence and the known degeneracy of the genetic code, a large number of different encoding nucleic acid sequences can be obtained. The degeneracy of the genetic code arises because almost  
20 all amino acids are encoded by different combinations of nucleotide triplets or "codons". Amino acids are encoded by codons as follows:

A=Ala=Alanine: codons GCA, GCC, GCG, GCU  
C=Cys=Cysteine: codons UGC, UGU  
D=Asp=Aspartic acid: codons GAC, GAU  
25 E=Glu=Glutamic acid: codons GAA, GAG  
F=Phe=Phenylalanine: codons UUC, UUU  
G=Gly=Glycine: codons GGA, GGC, GGG, GGU  
H=His=Histidine: codons CAC, CAU  
I=Ile=Isoleucine: codons AUA, AUC, AUU  
30 K=Lys=Lysine: codons AAA, AAG  
L=Leu=Leucine: codons UUA, UUG, CUA, CUC, CUG, CUU  
M=Met=Methionine: codon AUG  
N=Asn=Asparagine: codons AAC, AAU  
P=Pro=Proline: codons CCA, CCC, CCG, CCU  
35 Q=Gln=Glutamine: codons CAA, CAG

R=Arg=Arginine: codons AGA, AGG, CGA, CGC, CGG, CGU

S=Ser=Serine: codons AGC, AGU, UCA, UCC, UCG, UCU

T=Thr=Threonine: codons ACA, ACC, ACG, ACU

V=Val=Valine: codons GUA, GUC, GUG, GUU

5 W=Trp=Tryptophan: codon UGG

Y=Tyr=Tyrosine: codons UAC, UAU

Examples of nucleotide sequences encoding MCH-1R antagonist binding protein based on the human MCH-1R are provided by:

SEQ. ID. NO. 5: nucleotide MCH-1R(R141H);

10 SEQ. ID. NO. 6: nucleotide MCH-1R(R155A);

SEQ. ID. NO. 7: nucleotide MCH-1R(i2/MC4R); and

SEQ. ID. NO. 8: nucleotide MCH-1R( $\Delta$ 316/EGFP).

In different embodiments nucleic acid encoding a MCH-1R antagonist binding protein: (1) encode a protein having a sequence similarity of at least about 90%, preferably at least about 95% with either SEQ. ID. NOs. 1, 2, 3, 4, or a human MCH-1R with a deletion of about 37 amino acids; (2) encode a protein having a sequence with up to about 20 alterations from SEQ. ID. NOs. 1, 2, 3, 4, or a human MCH-1R with a deletion of about 37 amino acids; (3) the nucleic acid has a sequence similarity of at least about 90%, or at least about 95% with SEQ. ID. NO. 5, 6, 7, 8, or the human MCH-1R nucleic acid sequence with a deletion corresponding to about 37 C-terminal amino acids.

Sequence similarity for nucleic acid can be determined by FASTA. (Pearson 1990. *Methods in Enzymology* 183, 63-98, hereby incorporated by reference herein.) In one embodiment, sequence similarity is determined using the FASTA search program with the following parameters: MATRIX: BLOSUM50, GAP

25 PENALTIES: open=-12; residue=-2.

Nucleic acid having a desired sequence can be synthesized using chemical and biochemical techniques. Examples of chemical techniques are described in Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 1987-1998, and Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual*, 2<sup>nd</sup> Edition, Cold Spring Harbor Laboratory Press, 1989.

30

Biochemical nucleic acid synthesis techniques involve the use of a nucleic acid template and appropriate enzymes such as DNA and/or RNA polymerases. Examples of such techniques include *in vitro* amplification techniques such as PCR and transcription based amplification, and *in vivo* nucleic acid

35



replication. Examples of suitable techniques are provided by Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 1987-1998, Sambrook *et al.*, in *Molecular Cloning, A Laboratory Manual*, 2<sup>nd</sup> Edition, Cold Spring Harbor Laboratory Press, 1989, and Kacian, *et al.*, U.S. Patent No. 5,480,784.

5                   In an embodiment of the present invention, the MCH-1R antagonist binding protein is a purified polypeptide. A "purified polypeptide" represents at least 10% of the total protein present in a sample or preparation. In additional  
10                   embodiments, the purified polypeptide represents at least about 50%, at least about 75%, or at least about 95% of the total protein in a sample or preparation. Reference  
10                   to "purified polypeptide" does not require that the polypeptide has undergone any purification and may include, for example, chemically synthesized polypeptide that has not undergone any purification steps.

#### Recombinant Expression

15                   MCH-1R antagonist binding protein can be expressed from recombinant nucleic acid in a suitable host or in a test tube using a translation system. Recombinantly expressed MCH-1R antagonist binding protein are preferably used in assays to screen for compounds that bind to MCH-1R and modulate MCH-1R activity.

20                   Preferably, expression is achieved in a host cell using an expression vector. An expression vector is made up of recombinant nucleic acid encoding a polypeptide along with regulatory elements for proper transcription and processing. The regulatory elements that may be present include those naturally associated with the recombinant nucleic acid and exogenous regulatory elements not naturally  
25                   associated with the recombinant nucleic acid. Exogenous regulatory elements such as an exogenous promoter can be useful for expressing recombinant nucleic acid in a particular host.

                  Generally, the regulatory elements that are present in an expression vector include a transcriptional promoter, a ribosome binding site, a terminator, and  
30                   an optionally present operator. Another preferred element is a polyadenylation signal providing for processing in eukaryotic cells. Preferably, an expression vector also contains an origin of replication for autonomous replication in a host cell, a selectable marker, a limited number of useful restriction enzyme sites, and a potential for high copy number. Examples of expression vectors are cloning vectors, modified cloning  
35                   vectors, specifically designed plasmids and viruses.

Expression vectors providing suitable levels of polypeptide expression in different hosts are well known in the art. Mammalian expression vectors well known in the art include pcDNA3 (Invitrogen), pMC1neo (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), EBO-pSV2-neo (ATCC 37593), pBPV-1(8-2) (ATCC 37110), pdBPV-MMTneo(342-12) (ATCC 37224), pRSVgpt (ATCC 37199), pRSVneo (ATCC 37198), pSV2-dhfr (ATCC 37146), pUCTag (ATCC 37460), pCI-neo (Promega) and .lambda.ZD35 (ATCC 37565). Bacterial expression vectors well known in the art include pET11a (Novagen), lambda gt11 (Invitrogen), pcDNAII (Invitrogen), and pKK223-3 (Pharmacia). Fungal cell expression vectors well known in the art include pYES2 (Invitrogen) and Pichia expression vector (Invitrogen). Insect cell expression vectors well known in the art include Blue Bac III (Invitrogen).

Recombinant host cells may be prokaryotic or eukaryotic. Examples of recombinant host cells include the following: bacteria such as *E. coli*; fungal cells such as yeast; mammalian cells such as human, bovine, porcine, monkey and rodent; and insect cells such as *Drosophila* and silkworm derived cell lines. Commercially available mammalian cell lines include L cells L-M(TK.sup.-) (ATCC CCL 1.3), L cells L-M (ATCC CCL 1.2), 293 (ATCC CRL 1573), Raji (ATCC CCL 86), CV-1 (ATCC CCL 70), COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCC CCL 2), C127I (ATCC CRL 1616), BS-C-1 (ATCC CCL 26) and MRC-5 (ATCC CCL 171).

To enhance expression in a particular host it may be useful, for example, to modify the sequence provided in SEQ. ID. NOs. 5, 6, 7, or 8 to take into account codon usage of the host. Codon usage of different organisms are well known in the art. (See, Ausubel, *Current Protocols in Molecular Biology*, John Wiley, 1987-1998, Supplement 33 Appendix 1C.)

Expression vectors may be introduced into host cells using standard techniques. Examples of such techniques include transformation, transfection, lipofection, protoplast fusion, and electroporation.

Nucleic acid encoding an MCH-1R antagonist binding protein can be expressed in a cell without using of an expression vector by, for example, introducing a recombinant nucleic acid encoding the protein into the cell genome. Additionally, mRNA can be translated in various cell-free systems such as wheat germ extracts and reticulocyte extracts, as well as in cell based systems, such as frog oocytes.

Introduction of mRNA into cell based systems can be achieved, for example, by microinjection.

#### Functional Assays

5                   Evaluating the ability of a potential MCH-1R antagonist to modulate MCH-1R activity is facilitated through the use of an assay involving a functional MCH-1R and an MCH agonist. The use of an MCH agonist provides for MCH-1R activity.

10                   Recombinantly expressed MCH-1R can be used to facilitate determining receptor activity. For example, MCH-1R can be expressed by an expression vector in a cell line such as HEK 293, COS 7, or CHO, not normally expressing the receptor, wherein the same cell line without the expression vector or with an expression vector not encoding MCH-1R can act as a control.

15                   Functional assays can be performed using individual compounds or preparations containing different compounds. A preparation containing different compounds where one or more compounds affect MCH-1R activity can be divided into smaller groups of compounds to identify the compound(s) affecting MCH-1R activity.

#### 20                   Modulating MCH-1R Activity

                    MCH-1R antagonists have a variety of different uses including utility as a tool to further study MCH-1R activity and as an agent to achieve a beneficial effect in a patient. Beneficial effects of an MCH-1R antagonist include achieving one or more of the following in a patient: weight loss, cancer treatment (*e.g.*, colon or breast), pain reduction, diabetes treatment, stress reduction and sexual dysfunction treatment.

25                   A patient is a mammal, preferably a human. Reference to patient does not necessarily indicate the presence of a disease or disorder. The term patient includes subjects treated prophylactically and subjects afflicted with a disease or disorder.

30                   Excessive weight is a contributing factor to different diseases including hypertension, diabetes, dyslipidemias, cardiovascular disease, gall stones, osteoarthritis and certain forms of cancers. Bringing about a weight loss can be used, for example, to reduce the likelihood of such diseases and as part of a treatment for such diseases. Weight reduction can be achieved by, for example, one or more of the

35

following: reducing appetite, increasing metabolic rate, reducing fat intake and reducing carbohydrate craving.

Over weight patients include those having a body weight about 10% or more, 20% or more, 30% or more, or 50% or more, than the upper end of a "normal" weight range or Body Mass Index ("BMI"). "Normal" weight ranges are well known in the art and take into account factors such as a patient age, height, and body type.

BMI measures your height/weight ratio. It is determined by calculating weight in kilograms divided by the square of height in meters. The BMI "normal" range is 19-22.

MCH-1R modulating compounds can be provided in a kit. Such a kit typically contains an active compound in dosage forms for administration. A dosage form contains a sufficient amount of active compound such that a beneficial effect can be obtained when administered to a patient during regular intervals, such as 1 to 6 times a day, during the course of 1 or more days. Preferably, a kit contains instructions indicating the use of the dosage form for weight reduction (*e.g.*, to treat obesity or overweight) or stress reduction, and the amount of dosage form to be taken over a specified time period.

#### Dosing For Therapeutic Applications

Guidelines for pharmaceutical administration in general are provided in, for example, *Remington's Pharmaceutical Sciences 18<sup>th</sup> Edition*, Ed. Gennaro, Mack Publishing, 1990, and *Modern Pharmaceutics 2<sup>nd</sup> Edition*, Eds. Banker and Rhodes, Marcel Dekker, Inc., 1990, both of which are hereby incorporated by reference herein.

MCH-1R active compounds having appropriate functional groups can be prepared as acid or base salts. Pharmaceutically acceptable salts (in the form of water- or oil-soluble or dispersible products) include conventional non-toxic salts or the quaternary ammonium salts that are formed, *e.g.*, from inorganic or organic acids or bases. Examples of such salts include acid addition salts such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate,

pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, and undecanoate; and base salts such as ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine and lysine.

MCH-1R active compounds can be administered using different routes including oral, nasal, by injection, and transmucosally. Active ingredients to be administered orally as a suspension can be prepared according to techniques well known in the art of pharmaceutical formulation and may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners/flavoring agents. As immediate release tablets, these compositions may contain microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants.

When administered by nasal aerosol or inhalation, compositions can be prepared according to techniques well known in the art of pharmaceutical formulation. Such techniques can involve preparing solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, or other solubilizing or dispersing agents.

Routes of administration include intravenous (both bolus and infusion), intraperitoneal, subcutaneous, topical with or without occlusion, and intramuscular. Injectable solutions or suspensions known in the art include suitable non-toxic, parenterally-acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution and isotonic sodium chloride solution. Dispersing or wetting and suspending agents, include sterile, bland, fixed oils, such as synthetic mono- or diglycerides; and fatty acids, such as oleic acid.

Rectal administration in the form of suppositories include the use of a suitable non-irritating excipient, such as cocoa butter, synthetic glyceride esters or polyethylene glycols. These excipients are solid at ordinary temperatures, but liquidify and/or dissolve in the rectal cavity to release the drug.

Suitable dosing regimens for therapeutic applications can be designed taking into account factors well known in the art including age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound employed.

Optimal precision in achieving concentrations of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug. The daily dose for a patient is expected to be  
5 between 0.01 and 1,000 mg per adult patient per day.

#### EXAMPLES

Examples are provided below to further illustrate different features of the present invention. The examples also illustrate useful methodology for practicing  
10 the invention. These examples do not limit the claimed invention.

##### Example 1: Construction of MCH-1R(R141H), MCH-1R(R155A), MCH-1R( $\Delta$ 316/EGFP)

MCH-1R antagonist binding proteins were created by altering human  
15 MCH-1R. Alterations were generated using the QuikChange site-directed mutagenesis kit (Stratagene, La Jolla, CA) according to the manufacturers protocol. In brief: A template plasmid is denatured and mutant oligo-primers are annealed. Subsequently, using the non-strand-displacing action of PfuTurbo DNA polymerase primers are extended and incorporated in nicked circular strands. This step is repeated  
20 by thermal cycling. At the end of the reaction digestion of the methylated non-mutated parental DNA template is achieved by DpnI followed by transformation of the circular nicked DNA into supercompetent XL-1 Blue E. coli cells which repairs and amplifies the mutant plasmid.

The following combinations of mutant primers were used:

- 25 R141H+: 5'- CCATGGCCATTGACCACTACCTGGCCACTGTCC - 3' (SEQ. ID. NO. 9)  
R141H-: 5'- GGACAGTGGCCAGGTAGTGGTCAATGGCCATGG - 3' (SEQ. ID. NO. 10)
- 30 R155A+: 5'- CTCTTCCACGAAGTTCGCGAAGCCCTCTGTGGCC - 3' (SEQ. ID. NO. 11)  
R155A-: 5'- GGCCACAGAGGGCTTCGCGAACTTCGTGGAAGAG - 3' (SEQ. ID. NO. 12)

Δ316/EGFP+ :

5'-TTTGTGTACATCGTGCTCTGTGAGGTCGACGGTACCGCGGGCCCGGG- 3'  
(SEQ. ID NO. 13)

Δ316/EGFP- :

5 5'- CCCGGGCCCCGCGGTACCGTCGACCTCACAGAGCACGATGTACAC  
AAA- 3' (SEQ. ID. NO. 14)

The following templates were used:

MCH-1R(R141H): pcDNA3/MCH-1R  
10 MCH-1R(R155A): pcDNA3/MCH-1R  
MCH-1R(Δ316/EGFP): pEGFP-N3-MCH-1R

#### Example 2: Construction of MCH-1R(i2/MC4R)

MCH-1R(i2/MC4R) was created by PCR-based mutagenesis. The  
15 resulting MCH-1R antagonist binding protein contains the following amino acid  
sequence between TM3 and TM4: . . .DRYFTIFYALQYHNIMTVKRATLVICL  
(SEQ. ID NO. 15)... (the underlined sequence is the new sequence inserted in place of  
the original MCH-1R sequence.)

#### 20 Example 3: Analysis of Radioligand Binding

Membrane binding assays were performed using membrane  
preparations from transiently transfected HEK293-AEQ17 cells. HEK293-AEQ17  
cells (3-5 x 10<sup>6</sup> cells were plated in a T75 flask the day before transfection) were  
transiently transfected with plasmid DNA using LipofectAmine 2000 (Gibco BRL,  
25 Rockville, MD) according to the manufacturer's instructions. After two days  
membranes were prepared by hypotonic lysis, frozen in liquid nitrogen, and stored at  
- 80°C.

A scintillation proximity assay (SPA) was used to measure the specific  
binding of [<sup>125</sup>I]Phe<sup>13</sup>Tyr<sup>19</sup>-MCH (~2000 Ci/mmol; NEN Life Sciences, Boston, MA)  
30 to receptor containing membranes. SPA was carried out using wheat-germ agglutinin-  
polyvinyltoluene beads (Amersham Corp., Arlington Heights, IL), in 96-well  
OptiPlates (Packard, Meriden, CT). Each well contained 0.25 mg of SPA beads, 2 - 4  
μg of membrane protein, and 200 μl of binding buffer. Binding buffer contained 50  
mM Tris pH 7.4, 8 mM MgCl<sub>2</sub>, 12 % glycerol, 0.1 % BSA (Sigma, St. Louis, MO)

and protease inhibitors [4 µg/ml of leupeptin (Sigma, St. Louis, MO), 40 µg/ml of Bacitracin (Sigma, St. Louis, MO), 5 µg/ml of Aprotinin (Roche Molecular Biochem., Indianapolis, IN), and 100 µM AEBSF (Roche Molecular Biochem., Indianapolis, IN)].

5                    Assays were optimized with respect to membrane preparations: for HEK293-AEQ17/MCH-1R membranes, 1 µg of membranes per well yielded a > 6x specific binding window. Specific binding is defined as the difference between total binding and non-specific binding conducted in the presence of 500 nM unlabeled MCH. Beads were coated with membranes for 20 minutes and dispensed to the 96  
10 wells, various concentrations of test compounds in DMSO were added (final DMSO concentration 1 % - 2 %), then 25 nCi of [<sup>125</sup>I]Phe<sup>13</sup>Tyr<sup>19</sup>-MCH was added to the wells. After equilibrating at room temperature for 3 hours, the plates were read in a TopCount (Packard, Meriden, CT). IC<sub>50</sub> calculations were performed using Prism 3.0 (GraphPad Software, San Diego, CA).

15                    The results of the [<sup>125</sup>I]Phe<sup>13</sup>Tyr<sup>19</sup>-MCH agonist binding studies are shown in Figures 1 and 2. Figure 1 illustrates agonist binding to MCH-1R(R141H) and MCH-1R(R155A). Figure 2 illustrates agonist binding to MCH-1R(Δ316/EGFP).

#### Example 4: Functional Activation Analysis

20                    Functional activation of MCH-1R antagonist binding protein was measured in an aequorin assay. Proteins were introduced in the stable reporter cell line HEK293-AEQ17 in which mobilization of intracellular calcium can be detected by bioluminescence of jelly fish aequorin upon calcium binding.

25                    Bioluminescence was detected using a Luminoskan RT luminometer (Labsystems Inc., Gaithersburg, MD). HEK293-AEQ17 cells were maintained in D-MEM/high glucose medium (Life Technologies, Rockville, MD) supplemented with 10 % fetal bovine serum, 500 mg/ml G418, 25 mM Hepes at 37°C with 5 % CO<sub>2</sub> in a humidified atmosphere.

30                    HEK293-AEQ17 cells (3-5x 10<sup>6</sup> cells were plated in a T75 flask the day before transfection) were transiently transfected with MCH-1R antagonist binding protein plasmids using LipofectAmine 2000 (Gibco BRL, Rockville, MD) according to the manufacturer's instructions. After two days cells were incubated with the essential chromophore coelenterazine cp (10 µM; Molecular Probes, Eugene, OR) under reducing conditions (300 µM reduced glutathione in ECB buffer: 140 mM



NaCl, 20 mM KCl, 20 mM HEPES-NaOH pH 7.4, 5 mM glucose, 1 mM  $MgCl_2$ , 1 mM  $CaCl_2$ , 0.1 mg/ml BSA) to charge the apo-aequorin.

The cells were harvested, washed once in ECB medium and resuspended to 500,000 cells/ml. 100 ml of cell suspension (corresponding to  $5 \times 10^4$  cells) was then injected into a 96-well test plate, and the integrated light emission was recorded over 30 seconds, in 0.5 second units. 20  $\mu$ L of lysis buffer (0.1% final Triton X-100 concentration) was then injected and the integrated light emission recorded over 10 seconds, in 0.5 second units. The “fractional response” values for each well were calculated by taking the ratio of the integrated response to the initial challenge to the total integrated luminescence including the Triton X-100 lysis response.

### Example 5: Antagonist Binding

The ability of MCH-1R antagonist binding protein to bind an MCH-1R antagonist can be evaluated using standard techniques and techniques described herein. For example, the techniques described in Example 3 *supra*. can modified so that a labeled antagonist is employed.

Examples of MCH-1R antagonists are provided in U.S. Serial No. 60/310,928 (Attorney Docket NO. 20894PV), filed August 8, 2001, hereby  
20 incorporated by reference herein. Peptide antagonists include compounds having the structures ("\*" indicates cyclization (S-S)):

25 Ac-Gva-Cys-Met-Leu-Gly-Arg-Val-Tyr-Ava-Ava-Cys- NH<sub>2</sub> (SEQ. ID. NO. 16); and  
Ac-Gva-Cys-Met-Leu-D-Nle-Arg-Val-Tyr-Ava-Ava-Cys- NH<sub>2</sub> (SEQ. ID. NO. 17).

30 “Gva” refers to des-amino-arginine (also known as 5-guanidino-valeric acid). “Ava” refers to 5-aminovaleric acid. D-Nle refers to D-norleucine.

### Example 6: Sequence Information

Sequences for SEQ. ID. NO. 1-8 are provided below:

*MCH-1R(R141H) (SEQ. ID. NO. 1):*

MDLEASLLPTGPNASNTSDGPDNLTSAGSPPRTGSSISYINIIMPSVFGTICLLGIIG  
 NSTVIFAVVKKSKLHWCNNVPDIFIINLSVVDLLFLLGMPFMIHQLMGNGVWH  
 FGETMCTLITAMDANSQFTSTYILTAMAIDHYLATVHPISSTKFRKPSVATLVI  
 5 CLLWALSFSITPVWLYARLIPFPGGAVGCGIRLPNPDTDLYWFTLYQFFLAFA  
 LPFVVITAA YVRILQRMTSSVAPASQRSIRLRTKR VTRTAIAICLVFFVCWAPY  
 YVLQLTQLSISRPTLTFVYLYNAAISLGYANSCLNPFVYIVLCETFRKRLVLSV  
 KPAAQGQLRAVSNAQTADEERTESKGT

10 *MCH-1R(R155A) (SEQ. ID. NO. 2):*

MDLEASLLPTGPNASNTSDGPDNLTSAGSPPRTGSSISYINIIMPSVFGTICLLGIIG  
 NSTVIFAVVKKSKLHWCNNVPDIFIINLSVVDLLFLLGMPFMIHQLMGNGVWH  
 FGETMCTLITAMDANSQFTSTYILTAMAIDRYLATVHPISSTKFAKPSVATLVI  
 CLLWALSFSITPVWLYARLIPFPGGAVGCGIRLPNPDTDLYWFTLYQFFLAFA  
 15 LPFVVITAA YVRILQRMTSSVAPASQRSIRLRTKR VTRTAIAICLVFFVCWAPY  
 YVLQLTQLSISRPTLTFVYLYNAAISLGYANSCLNPFVYIVLCETFRKRLVLSV  
 KPAAQGQLRAVSNAQTADEERTESKGT

## MCH-1R(i2/MC4R) (SEQ. ID. NO. 3)

20 MDLEASLLPTGPNASNTSDGPDNLTSAGSPPRTGSSISYINIIMPSVFGTICLLGIIG  
 NSTVIFAVVKKSKLHWCNNVPDIFIINLSVVDLLFLLGMPFMIHQLMGNGVWH  
 FGETMCTLITAMDANSQFTSTYILTAMAIDRY **FTIFYALQYHNIMTVKR**  
 ATL VICLLWALSFSITPVWLYARLIPFPGGAVGCGIRLPNPDTDLYWFTLYQFF  
 LAFALPFVVITAA YVRILQRMTSSVAPASQRSIRLRTKR VTRTAIAICLVFFVC  
 25 WAPYYVLQLTQLSISRPTLTFVYLYNAAISLGYANSCLNPFVYIVLCETFRKRL  
 VLSVKPAAQGQLRAVSNAQTADEERTESKGT

*MCH-1R(Δ316/EGFP) (SEQ. ID. NO. 4)*

MDLEASLLPTGPNASNTSDGPDNLTSAGSPPRTGSSISYINIIMPSVFGTICLLGIIG  
 30 NSTVIFAVVKKSKLHWCNNVPDIFIINLSVVDLLFLLGMPFMIHQLMGNGVWH  
 FGETMCTLITAMDANSQFTSTYILTAMAIDRYLATVHPISSTKFRKPSVATLVI  
 CLLWALSFSITPVWLYARLIPFPGGAVGCGIRLPNPDTDLYWFTLYQFFLAFA  
 LPFVVITAA YVRILQRMTSSVAPASQRSIRLRTKR VTRTAIAICLVFFVCWAPY  
 YVLQLTQLSISRPTLTFVYLYNAAISLGYANSCLNPFVYIVLCEVDGTAGPGSI  
 35 ATMVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTT

GKLPVPWPTLVTTLT YGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKD  
 DGNKYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMA  
 DKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSAL  
 SKDPNEKRDHMLLEFVTAAGITLGMDELYK

5

*MCH-1R(R141H)* (SEQ. ID. NO. 5): Start and stop codons as well as mutant nucleotide are highlighted.

ATGGACCTGGAAGCCTCGCTGCTGCCCCACTGGTCCCAATGCCAGCAACAC  
 CTCTGATGGCCCCGATAACCTCACTTCGGCAGGATCACCTCCTCGCACGG  
 10 GGAGCATCTCCTACATCAACATCATCATGCCTTCGGTGTTCGGCACCATCT  
 GCCTCCTGGGCATCATCGGGAACCTCCACGGTCATCTTCGCGGTCTGTGAAG  
 AAGTCCAAGCTGCACTGGTGCAACAACGTCCCCGACATCTTCATCATCAA  
 CCTCTCGGTAGTAGATCTCCTCTTTCTCCTGGGCATGCCCTTCATGATCCA  
 CCAGCTCATGGGCAATGGGGTGTGGCACTTTGGGGAGACCATGTGCACCC  
 15 TCATCACGGCCATGGATGCCAATAGTCAGTTCACCAGCACCTACATCCTG  
 ACCGCCATGGCCATTGACCACTACCTGGCCACTGTCCACCCCATCTCTTCC  
 ACGAAGTTCCGGAAGCCCTCTGTGGCCACCCTGGTGATCTGCCTCCTGTGG  
 GCCCTCTCCTTCATCAGCATCACCCCTGTGTGGCTGTATGCCAGACTCATC  
 CCCTTCCCAGGAGGTGCAGTGGGCTGCGGCATACGCCTGCCCAACCCAGA  
 20 CACTGACCTCTACTGGTTCACCCTGTACCAGTTTTTCCTGGCCTTTGCCCTG  
 CCTTTTGTGGTCATCACAGCCGCATACGTGAGGATCCTGCAGCGCATGAC  
 GTCCTCAGTGGCCCCCGCCTCCAGCGCAGCATCCGGCTGCGGACAAAGA  
 GGGTGACCCGCACAGCCATCGCCATCTGTCTGGTCTTCTTTGTGTGCTGGG  
 CACCCTACTATGTGCTACAGCTGACCCAGTTGTCCATCAGCCGCCCCGACCC  
 25 TCACCTTTGTCTACTTATACAATGCGGCCATCAGCTTGGGCTATGCCAACA  
 GCTGCCTCAACCCCTTTGTGTACATCGTGCTCTGTGAGACGTTCCGCAAAC  
 GCTTGGTCCTGTCCGTGAAGCCTGCAGCCCAGGGGCAGCTTCGCGCTGTC  
 AGCAACGCTCAGACGGCTGACGAGGAGAGGACAGAAAGCAAAGGCACCT  
 GATAC

30

*MCH-1R(R141A)* (SEQ. ID. NO. 6): Start and stop codons as well as mutant nucleotide are highlighted. Nucleic acid sequence (start and stop codons as well as mutant nucleotide are highlighted).

ATGGACCTGGAAGCCTCGCTGCTGCCCCACTGGTCCCAATGCCAGCAACAC  
 35 CTCTGATGGCCCCGATAACCTCACTTCGGCAGGATCACCTCCTCGCACGG

GGAGCATCTCCTACATCAACATCATCATGCCTTCGGTGTTCCGGCACCATCT  
 GCCTCCTGGGCATCATCGGGAAGTCCACGGTCATCTTCGCGGTCGTGAAG  
 AAGTCCAAGCTGCACTGGTGCAACAACGTCCCCGACATCTTCATCATCAA  
 CCTCTCGGTAGTAGATCTCCTCTTTCTCCTGGGCATGCCCTTCATGATCCA  
 5 CCAGCTCATGGGCAATGGGGTGTGGCACTTTGGGGAGACCATGTGCACCC  
 TCATCACGGCCATGGATGCCAATAGTCAGTTCACCAGCACCTACATCCTG  
 ACCGCCATGGCCATTGACCGCTACCTGGCCACTGTCCACCCCATCTCTTCC  
 ACGAAGTTCGCGAAGCCCTCTGTGGCCACCCTGGTGATCTGCCTCCTGTG  
 GGCCCTCTCCTTCATCAGCATCACCCCTGTGTGGCTGTATGCCAGACTCAT  
 10 CCCCTTCCCAGGAGGTGCAGTGGGCTGCGGCATACGCCTGCCCAACCCAG  
 AACTGACCTCTACTGGTTCACCCTGTACCAGTTTTTCCTGGCCTTTGCCCT  
 GCCTTTTGTGGTCATCACAGCCGCATACGTGAGGATCCTGCAGCGCATGA  
 CGTCCTCAGTGGCCCCCGCCTCCCAGCGCAGCATCCGGCTGCGGACAAAG  
 AGGGTGACCCGCACAGCCATCGCCATCTGTCTGGTCTTCTTTGTGTGCTGG  
 15 GCACCCTACTATGTGCTACAGCTGACCCAGTTGTCCATCAGCCGCCCCGAC  
 CCTCACCTTTGTCTACTTATAACAATGCGGCCATCAGCTTGGGCTATGCCAA  
 CAGCTGCCTCAACCCCTTTGTGTACATCGTGCTCTGTGAGACGTTCCGCAA  
 ACGCTTGGTCCTGTGCGTGAAGCCTGCAGCCCAGGGGCAGCTTCGCGCTG  
 TCAGCAACGCTCAGACGGCTGACGAGGAGAGGACAGAAAGCAAAGGCAC  
 20 CTGATAC

*MCH-1R(i2/MC4R)* (SEQ. ID. NO. 7): Start and stop codons as well as mutant nucleotide are highlighted.

ATGGACCTGGAAGCCTCGCTGCTGCCCCACTGGTCCCAATGCCAGCAACAC  
 25 CTCTGATGGCCCCGATAACCTCACTTCGGCAGGATCACCTCCTCGCACGG  
 GGAGCATCTCCTACATCAACATCATCATGCCTTCGGTGTTCCGGCACCATCT  
 GCCTCCTGGGCATCATCGGGAAGTCCACGGTCATCTTCGCGGTCGTGAAG  
 AAGTCCAAGCTGCACTGGTGCAACAACGTCCCCGACATCTTCATCATCAA  
 CCTCTCGGTAGTAGATCTCCTCTTTCTCCTGGGCATGCCCTTCATGATCCA  
 30 CCAGCTCATGGGCAATGGGGTGTGGCACTTTGGGGAGACCATGTGCACCC  
 TCATCACGGCCATGGATGCCAATAGTCAGTTCACCAGCACCTACATCCTG  
 ACCGCCATGGCCATTGACCGCTACTTTACTATCTTCTATGCTCTCCAGTACC  
 ATAACATTATGACAGTTAAGCGGGCCACCCTGGTGATCTGCCTCCTGTGGG  
 CCCTCTCCTTCATCAGCATCACCCCTGTGTGGCTGTATGCCAGACTCATCC  
 35 CCTTCCCAGGAGGTGCAGTGGGCTGCGGCATACGCCTGCCCAACCCAGAC

ACTGACCTCTACTGGTTACCCCTGTACCAGTTTTTCCTGGCCTTTGCCCTGC  
 CTTTTGTGGTCATCACAGCCGCATACGTGAGGATCCTGCAGCGCATGACG  
 TCCTCAGTGGCCCCCGCCTCCCAGCGCAGCATCCGGCTGCGGACAAAGAG  
 GGTGACCCGCACAGCCATCGCCATCTGTCTGGTCTTCTTTGTGTGCTGGGC  
 5 ACCCTACTATGTGCTACAGCTGACCCAGTTGTCCATCAGCCGCCCCGACCCT  
 CACCTTTGTCTACTTATACAATGCGGCCATCAGCTTGGGCTATGCCAACAG  
 CTGCCTCAACCCCTTTGTGTACATCGTGCTCTGTGAGACGTTCCGCAAACG  
 CTTGGTCTGTGCGGTGAAGCCTGCAGCCCAGGGGCAGCTTCGCGCTGTCA  
 GCAACGCTCAGACGGCTGACGAGGAGAGGACAGAAAGCAAAGGCACCTG  
 10 ATAC

*MCH-1R(Δ316/EGFP) (SEQ. ID. NO. 8):* Start codon and start and stop codons for  
 MCH-1R and EGFP, respectively, are highlighted. A 12 amino acid linker sequence  
 is denoted in lower case.

15 ATGGACCTGGAAGCCTCGCTGCTGCCCCTGGTCCCAATGCCAGCAACAC  
 CTCTGATGGCCCCGATAACCTCACTTCGGCAGGATCACCTCCTCGCACGG  
 GGAGCATCTCCTACATCAACATCATCATGCCTTCGGTGTTCCGGCACCATCT  
 GCCTCCTGGGCATCATCGGGAACCTCCACGGTCATCTTCGCGGTCTGTGAAG  
 AAGTCCAAGCTGCACTGGTGCAACAACGTCCCCGACATCTTCATCATCAA  
 20 CCTCTCGGTAGTAGATCTCCTCTTTCTCCTGGGCATGCCCTTCATGATCCA  
 CCAGCTCATGGGCAATGGGGTGTGGCACTTTGGGGAGACCATGTGCACCC  
 TCATCACGGCCATGGATGCCAATAGTCAGTTCACCAGCACCTACATCCTG  
 ACCGCCATGGCCATTGACCGCTACCTGGCCACTGTCCACCCCATCTCTTCC  
 ACGAAGTTCCGGAAGCCCTCTGTGGCCACCCTGGTGATCTGCCTCCTGTGG  
 25 GCCCTCTCCTTCATCAGCATCACCCCTGTGTGGCTGTATGCCAGACTCATC  
 CCCTTCCCAGGAGGTGCAGTGGGCTGCGGCATACGCCTGCCCAACCCAGA  
 CACTGACCTCTACTGGTTACCCCTGTACCAGTTTTTCCTGGCCTTTGCCCTG  
 CCTTTTGTGGTCATCACAGCCGCATACGTGAGGATCCTGCAGCGCATGAC  
 GTCCTCAGTGGCCCCCGCCTCCCAGCGCAGCATCCGGCTGCGGACAAAGA  
 30 GGGTGACCCGCACAGCCATCGCCATCTGTCTGGTCTTCTTTGTGTGCTGGG  
 CACCCTACTATGTGCTACAGCTGACCCAGTTGTCCATCAGCCGCCCCGACCC  
 TCACCTTTGTCTACTTATACAATGCGGCCATCAGCTTGGGCTATGCCAACAA  
 GCTGCCTCAACCCCTTTGTGTACATCGTGCTCTGTGAGgtcgacggtaccgaggcc  
 gggatccatgccaccATGGTG AGCAAGGGCG AGGAGCTGTT CACCGGGGTG  
 35 GTGCCCATCC TGGTCGAGCT GGACGGCGAC GTAAACGGCC

ACAAGTTCAG CGTGTCCGGC GAGGGCGAGG GCGATGCCAC  
CTACGGCAAG CTGACCCTGA AGTTCATCTG CACCACCGGC  
AAGCTGCCCCG TGCCCTGGCC CACCCTCGTG ACCACCCTGA  
CCTACGGCGT GCAGTGCTTC AGCCGCTACC CCGACCACAT  
5 GAAGCAGCAC GACTTCTTCA AGTCCGCCAT GCCC GAAGGC  
TACGTCCAGG AGCGCACCAT CTTCTTCAAG GACGACGGCA  
ACTACAAGAC CCGCGCCGAG GTGAAGTTCG AGGGCGACAC  
CCTGGTGAAC CGCATCGAGC TGAAGGGCAT CGACTTCAAG  
GAGGACGGCA ACATCCTGGG GCACAAGCTG GAGTACAAC  
10 ACAACAGCCA CAACGTCTAT ATCATGGCCG ACAAGCAGAA  
GAACGGCATC AAGGTGAACT TCAAGATCCG CCACAACATC  
GAGGACGGCA GCGTGCAGCT CGCCGACCAC TACCAGCAGA  
ACACCCCCAT CCGCGACGGC CCCGTGCTGC TGCCCGACAA  
CCACTACCTG AGCACCAGT CCGCCCTGAG CAAAGACCCC  
15 AACGAGAAGC GCGATCACAT GGTCTGCTG GAGTTCGTGA  
CCGCCGCCGG GATCACTCTC GGCATGGACG AGCTGTACAA GTAA

Other embodiments are within the following claims. While several  
20 embodiments have been shown and described, various modifications may be made  
without departing from the spirit and scope of the present invention.

## WHAT IS CLAIMED IS:

1. A melanin-concentrating hormone receptor type 1 (MCH-1R) antagonist binding protein selected from the group consisting of:
  - 5 a) a first MCH-1R antagonist binding protein, wherein said first MCH-1R antagonist binding protein has one or more alterations in the second intracellular loop region that render MCH-1R substantially inactive to agonist activation; and
  - 10 b) a second MCH-1R antagonist binding protein, wherein said second MCH-1R antagonist binding protein has one or more alterations in the C-terminal region that render MCH-1R substantially inactive to agonist activation.
- 15 2. The MCH-1R antagonist binding protein of claim 1, wherein said MCH-1R antagonist binding protein is said first MCH-1R antagonist binding protein.
- 20 3. The MCH-1R antagonist binding protein of claim 2, wherein said MCH-1R antagonist binding protein consists of the amino acid sequence of SEQ ID NO: 1.
- 25 4. The MCH-1R antagonist binding protein of claim 2, wherein said MCH-1R antagonist binding protein consists of the amino acid sequence of SEQ ID NO: 2.
- 30 5. The MCH-1R antagonist binding protein of claim 2, wherein said MCH-1R antagonist binding protein consists of the amino acid sequence of SEQ ID NO: 3.
- 35 6. The MCH-1R antagonist binding protein of claim 1, wherein said MCH-1R antagonist binding protein is said second MCH-1R antagonist binding protein.
7. The MCH-1R antagonist binding protein of claim 6, wherein said MCH-1R antagonist binding protein consists of the amino acid sequence of SEQ ID NO: 4.

8. A nucleic acid comprising a nucleotide sequence encoding the MCH-1R antagonist binding protein of any one of claims 1-7.
- 5 9. A nucleic acid comprising a nucleotide sequence selected from the group consisting of: SEQ ID NO: 5, SEQ ID NO: 6., SEQ ID NO: 7, and SEQ ID NO: 8.
- 10 10. The nucleic acid of claim 9, wherein said nucleotide sequence consists of SEQ ID NO: 5.
11. The nucleic acid of claim 9, wherein said nucleotide sequence consists of SEQ ID NO: 6.
- 15 12. The nucleic acid of claim 9, wherein said nucleotide sequence consists of SEQ ID NO: 7.
13. The nucleic acid of claim 9, wherein said nucleotide sequence consists of SEQ ID NO: 8.
- 20 14. The nucleic acid of any one of claims 8-13, wherein said nucleic acid is an expression vector.
15. A recombinant cell comprising the expression vector of claim 25 14, wherein said nucleotide sequence is functionally coupled to a promoter recognized by said cell.
16. A method of screening for a compound able to bind MCH-1R comprising the steps of:
- 30 a) contacting the MCH-1R antagonist binding protein of any one of claims 1-7 with a said compound; and
- b) measuring the ability of said compound to bind to said MCH-1R antagonist binding protein.



17. A method of preparing a MCH-1R antagonist binding protein comprising the step of growing the recombinant cell of claim 15 under conditions wherein said protein is expressed from said expression vector.

5

1/6




FIG. 1

2/6




FIG. 2

3/6




FIG. 3

4/6




FIG. 4

5/6




FIG. 5

6/6




FIG. 6

## SEQUENCE LISTING

&lt;110&gt; Merck &amp; Co., Inc.

<120> MELANIN-CONCENTRATING HORMONE RECEPTOR  
ANTAGONIST BINDING PROTEIN

&lt;130&gt; 20782 PCT

&lt;150&gt; 60/325,129

&lt;151&gt; 2001-09-26

&lt;160&gt; 17

&lt;170&gt; FastSEQ for Windows Version 4.0

&lt;210&gt; 1

&lt;211&gt; 353

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; MCH-1R antagonist binding protein

&lt;400&gt; 1

```

Met Asp Leu Glu Ala Ser Leu Leu Pro Thr Gly Pro Asn Ala Ser Asn
 1          5          10          15
Thr Ser Asp Gly Pro Asp Asn Leu Thr Ser Ala Gly Ser Pro Pro Arg
      20          25          30
Thr Gly Ser Ile Ser Tyr Ile Asn Ile Ile Met Pro Ser Val Phe Gly
      35          40          45
Thr Ile Cys Leu Leu Gly Ile Ile Gly Asn Ser Thr Val Ile Phe Ala
 50          55          60
Val Val Lys Lys Ser Lys Leu His Trp Cys Asn Asn Val Pro Asp Ile
 65          70          75          80
Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu Phe Leu Leu Gly Met
      85          90          95
Pro Phe Met Ile His Gln Leu Met Gly Asn Gly Val Trp His Phe Gly
      100          105          110
Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp Ala Asn Ser Gln Phe
      115          120          125
Thr Ser Thr Tyr Ile Leu Thr Ala Met Ala Ile Asp His Tyr Leu Ala
      130          135          140
Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg Lys Pro Ser Val Ala
 145          150          155          160
Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser Phe Ile Ser Ile Thr
      165          170          175
Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe Pro Gly Gly Ala Val
      180          185          190
Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr Asp Leu Tyr Trp Phe
      195          200          205
Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu Pro Phe Val Val Ile
      210          215          220
Thr Ala Ala Tyr Val Arg Ile Leu Gln Arg Met Thr Ser Ser Val Ala
 225          230          235          240

```



<400>	2															
Met	Asp	Leu	Glu	Ala	Ser	Leu	Leu	Pro	Thr	Gly	Pro	Asn	Ala	Ser	Asn	
1				5					10					15		
Thr	Ser	Asp	Gly	Pro	Asp	Asn	Leu	Thr	Ser	Ala	Gly	Ser	Pro	Pro	Arg	
			20					25					30			
Thr	Gly	Ser	Ile	Ser	Tyr	Ile	Asn	Ile	Ile	Met	Pro	Ser	Val	Phe	Gly	
		35					40					45				
Thr	Ile	Cys	Leu	Leu	Gly	Ile	Ile	Gly	Asn	Ser	Thr	Val	Ile	Phe	Ala	
		50				55					60					
Val	Val	Lys	Lys	Ser	Lys	Leu	His	Trp	Cys	Asn	Val	Pro	Asp	Ile		
65					70				75					80		
Phe	Ile	Ile	Asn	Leu	Ser	Val	Val	Asp	Leu	Leu	Phe	Leu	Leu	Gly	Met	
				85					90					95		
Pro	Phe	Met	Ile	His	Gln	Leu	Met	Gly	Asn	Gly	Val	Trp	His	Phe	Gly	
			100					105					110			
Glu	Thr	Met	Cys	Thr	Leu	Ile	Thr	Ala	Met	Asp	Ala	Asn	Ser	Gln	Phe	
		115					120					125				
Thr	Ser	Thr	Tyr	Ile	Leu	Thr	Ala	Met	Ala	Ile	Asp	Arg	Tyr	Leu	Ala	
		130				135					140					
Thr	Val	His	Pro	Ile	Ser	Ser	Thr	Lys	Phe	Ala	Lys	Pro	Ser	Val	Ala	
145					150				155						160	
Thr	Leu	Val	Ile	Cys	Leu	Leu	Trp	Ala	Leu	Ser	Phe	Ile	Ser	Ile	Thr	
				165					170					175		
Pro	Val	Trp	Leu	Tyr	Ala	Arg	Leu	Ile	Pro	Phe	Pro	Gly	Gly	Ala	Val	
			180					185					190			
Gly	Cys	Gly	Ile	Arg	Leu	Pro	Asn	Pro	Asp	Thr	Asp	Leu	Tyr	Trp	Phe	
		195					200					205				
Thr	Leu	Tyr	Gln	Phe	Phe	Leu	Ala	Phe	Ala	Leu	Pro	Phe	Val	Val	Ile	
		210				215					220					
Thr	Ala	Ala	Tyr	Val	Arg	Ile	Leu	Gln	Arg	Met	Thr	Ser	Ser	Val	Ala	
225					230					235					240	

Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr Lys Arg Val Thr Arg  
                                   245                                  250                                  255  
 Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val Cys Trp Ala Pro Tyr  
                                   260                                  265                                  270  
 Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser Arg Pro Thr Leu Thr  
                                   275                                  280                                  285  
 Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu Gly Tyr Ala Asn Ser  
                                   290                                  295                                  300  
 Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys Glu Thr Phe Arg Lys  
 305                                  310                                  315                                  320  
 Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln Gly Gln Leu Arg Ala  
                                   325                                  330                                  335  
 Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg Thr Glu Ser Lys Gly  
                                   340                                  345                                  350  
 Thr

<210> 3

<211> 353

<212> PRT

<213> Artificial Sequence

<220>

<223> MCH-1R antagonist binding protein

<400> 3

Met Asp Leu Glu Ala Ser Leu Leu Pro Thr Gly Pro Asn Ala Ser Asn  
 1                                  5                                  10                                  15  
 Thr Ser Asp Gly Pro Asp Asn Leu Thr Ser Ala Gly Ser Pro Pro Arg  
                                   20                                  25                                  30  
 Thr Gly Ser Ile Ser Tyr Ile Asn Ile Ile Met Pro Ser Val Phe Gly  
                                   35                                  40                                  45  
 Thr Ile Cys Leu Leu Gly Ile Ile Gly Asn Ser Thr Val Ile Phe Ala  
                                   50                                  55                                  60  
 Val Val Lys Lys Ser Lys Leu His Trp Cys Asn Asn Val Pro Asp Ile  
 65                                  70                                  75                                  80  
 Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu Phe Leu Leu Gly Met  
                                   85                                  90                                  95  
 Pro Phe Met Ile His Gln Leu Met Gly Asn Gly Val Trp His Phe Gly  
                                   100                                  105                                  110  
 Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp Ala Asn Ser Gln Phe  
                                   115                                  120                                  125  
 Thr Ser Thr Tyr Ile Leu Thr Ala Met Ala Ile Asp Arg Tyr Phe Thr  
                                   130                                  135                                  140  
 Ile Phe Tyr Ala Leu Gln Tyr His Asn Ile Met Thr Val Lys Arg Ala  
 145                                  150                                  155                                  160  
 Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser Phe Ile Ser Ile Thr  
                                   165                                  170                                  175  
 Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe Pro Gly Gly Ala Val  
                                   180                                  185                                  190  
 Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr Asp Leu Tyr Trp Phe  
                                   195                                  200                                  205  
 Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu Pro Phe Val Val Ile  
                                   210                                  215                                  220  
 Thr Ala Ala Tyr Val Arg Ile Leu Gln Arg Met Thr Ser Ser Val Ala  
 225                                  230                                  235                                  240

```

Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr Lys Arg Val Thr Arg
                245                250                255
Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val Cys Trp Ala Pro Tyr
                260                265                270
Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser Arg Pro Thr Leu Thr
                275                280                285
Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu Gly Tyr Ala Asn Ser
                290                295                300
Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys Glu Thr Phe Arg Lys
305                310                315                320
Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln Gly Gln Leu Arg Ala
                325                330                335
Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg Thr Glu Ser Lys Gly
                340                345                350

```

Thr

&lt;210&gt; 4

&lt;211&gt; 567

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; MCH-1R antagonist binding protein

&lt;400&gt; 4

```

Met Asp Leu Glu Ala Ser Leu Leu Pro Thr Gly Pro Asn Ala Ser Asn
1          5          10          15
Thr Ser Asp Gly Pro Asp Asn Leu Thr Ser Ala Gly Ser Pro Pro Arg
          20          25          30
Thr Gly Ser Ile Ser Tyr Ile Asn Ile Ile Met Pro Ser Val Phe Gly
          35          40          45
Thr Ile Cys Leu Leu Gly Ile Ile Gly Asn Ser Thr Val Ile Phe Ala
50          55          60
Val Val Lys Lys Ser Lys Leu His Trp Cys Asn Asn Val Pro Asp Ile
65          70          75          80
Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu Phe Leu Leu Gly Met
          85          90          95
Pro Phe Met Ile His Gln Leu Met Gly Asn Gly Val Trp His Phe Gly
          100         105         110
Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp Ala Asn Ser Gln Phe
          115         120         125
Thr Ser Thr Tyr Ile Leu Thr Ala Met Ala Ile Asp Arg Tyr Leu Ala
          130         135         140
Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg Lys Pro Ser Val Ala
145         150         155         160
Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser Phe Ile Ser Ile Thr
          165         170         175
Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe Pro Gly Gly Ala Val
          180         185         190
Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr Asp Leu Tyr Trp Phe
          195         200         205
Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu Pro Phe Val Val Ile
210         215         220
Thr Ala Ala Tyr Val Arg Ile Leu Gln Arg Met Thr Ser Ser Val Ala
225         230         235         240

```

```

Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr Lys Arg Val Thr Arg
                245                250                255
Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val Cys Trp Ala Pro Tyr
                260                265                270
Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser Arg Pro Thr Leu Thr
                275                280                285
Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu Gly Tyr Ala Asn Ser
                290                295                300
Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys Glu Val Asp Gly Thr
305                310                315                320
Ala Gly Pro Gly Ser Ile Ala Thr Met Val Ser Lys Gly Glu Glu Leu
                325                330                335
Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn
                340                345                350
Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr
                355                360                365
Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val
370                375                380
Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe
385                390                395                400
Ser Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala
                405                410                415
Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp
                420                425                430
Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu
435                440                445
Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn
450                455                460
Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr
465                470                475                480
Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile
                485                490                495
Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln
                500                505                510
Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His
515                520                525
Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg
530                535                540
Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu
545                550                555                560
Gly Met Asp Glu Leu Tyr Lys
                565

```

&lt;210&gt; 5

&lt;211&gt; 1065

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; MCH-1R antagonist binding protein

&lt;400&gt; 5

```

atggacctgg aagcctcgct gctgccact ggtcccaatg ccagcaacac ctctgatggc    60
cccgataacc tcaattcggc aggatcacct cctcgcacgg ggagcatctc ctacatcaac    120
atcatcatgc cttcgggtgtt cggcaccatc tgcctcctgg gcatcatcgg gaactccacg    180

```

gtcatcttcg	cggtcgtgaa	gaagtccaag	ctgcactggg	gcaacaacgt	ccccgacatc	240
ttcatcatca	acctctcggt	agtagatctc	ctctttctcc	tgggcatgcc	cttcatgata	300
caccagctca	tgggcaatgg	ggtgtggcac	tttggggaga	ccatgtgcac	cctcatcacg	360
gccatggatg	ccaatagtca	gttcaccagc	acctacatcc	tgaccgccat	ggccattgac	420
cactacctgg	ccactgtcca	ccccatctct	tccacgaagt	tccggaagcc	ctctgtggcc	480
accttgggtg	tctgcctcct	gtgggcccct	tccttcatca	gcatcacccc	tgtgtggctg	540
tatgccagac	tcatcccctt	cccaggaggt	gcagtgggct	gcggcatacg	cctgccccaa	600
ccagacactg	acctctactg	gttcaccctg	taccagtttt	tcctggcctt	tgccctgcct	660
tttgtgggtc	tcacagccgc	atacgtgagg	atcctgcagc	gcatgacgtc	ctcagtggcc	720
cccgcctccc	agcgcagcat	ccggctgcgg	acaaagaggg	tgacccgcac	agccatcgcc	780
atctgtctgg	tcttctttgt	gtgctgggca	ccctactatg	tgctacagct	gaccagttg	840
tccatcagcc	gcccagacct	cacctttgtc	tacttatata	atgcggccat	cagcttgggc	900
tatgccaaca	gctgcctcaa	cccctttgtg	tacatcgtgc	tctgtgagac	gttccgcaaa	960
cgcttgggtc	tgtcgggtgaa	gcctgcagcc	caggggcagc	ttcgcgctgt	cagcaacgct	1020
cagacggctg	acgaggagag	gacagaaagc	aaaggcacct	gatac		1065

&lt;210&gt; 6

&lt;211&gt; 1065

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; MCH-1R antagonist binding protein

&lt;400&gt; 6

atggacctgg	aagcctcgct	gctgcccact	ggccccaatg	ccagcaaacac	ctctgatggc	60
cccgataacc	tcacttcggc	aggatcacct	cctcgcacgg	ggagcatctc	ctacatcaac	120
atcatcatgc	cttcgggtgt	cggcaccatc	tgcctcctgg	gcatcatcgg	gaactccacg	180
gtcatcttcg	cggtcgtgaa	gaagtccaag	ctgcactggg	gcaacaacgt	ccccgacatc	240
ttcatcatca	acctctcggt	agtagatctc	ctctttctcc	tgggcatgcc	cttcatgata	300
caccagctca	tgggcaatgg	ggtgtggcac	tttggggaga	ccatgtgcac	cctcatcacg	360
gccatggatg	ccaatagtca	gttcaccagc	acctacatcc	tgaccgccat	ggccattgac	420
cgctacctgg	ccactgtcca	ccccatctct	tccacgaagt	tccggaagcc	ctctgtggcc	480
accttgggtg	tctgcctcct	gtgggcccct	tccttcatca	gcatcacccc	tgtgtggctg	540
tatgccagac	tcatcccctt	cccaggaggt	gcagtgggct	gcggcatacg	cctgccccaa	600
ccagacactg	acctctactg	gttcaccctg	taccagtttt	tcctggcctt	tgccctgcct	660
tttgtgggtc	tcacagccgc	atacgtgagg	atcctgcagc	gcatgacgtc	ctcagtggcc	720
cccgcctccc	agcgcagcat	ccggctgcgg	acaaagaggg	tgacccgcac	agccatcgcc	780
atctgtctgg	tcttctttgt	gtgctgggca	ccctactatg	tgctacagct	gaccagttg	840
tccatcagcc	gcccagacct	cacctttgtc	tacttatata	atgcggccat	cagcttgggc	900
tatgccaaca	gctgcctcaa	cccctttgtg	tacatcgtgc	tctgtgagac	gttccgcaaa	960
cgcttgggtc	tgtcgggtgaa	gcctgcagcc	caggggcagc	ttcgcgctgt	cagcaacgct	1020
cagacggctg	acgaggagag	gacagaaagc	aaaggcacct	gatac		1065

&lt;210&gt; 7

&lt;211&gt; 1065

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; MCH-1R antagonist binding protein

&lt;400&gt; 7

atggacctgg	aagcctcgct	gctgcccact	ggccccaatg	ccagcaaacac	ctctgatggc	60
cccgataacc	tcacttcggc	aggatcacct	cctcgcacgg	ggagcatctc	ctacatcaac	120

atcatcatgc	cttcggtggt	cggcaccatc	tgctcctcgtg	gcatcatcgg	gaactccacg	180
gtcatcttcg	cggtcgtgaa	gaagtccaag	ctgcactggg	gcaacaacgt	ccccgacatc	240
ttcatcatca	acctctcggg	agtagatctc	ctctttctcc	tgggcatgcc	cttcatgatc	300
caccagctca	tgggcaatgg	ggtgtggcac	tttggggaga	ccatgtgcac	cctcatcacg	360
gccatggatg	ccaatagtca	gttcaccagc	acctacatcc	tgaccgccat	ggccattgac	420
cgctacttta	ctatcttcta	tgctctccag	taccataaca	ttatgacagt	taagcggggc	480
accctgggtga	tctgcctcct	gtgggcccctc	tccttcatca	gcatcacccc	tgtgtggctg	540
tatgccagac	tcattcccctt	cccaggaggt	gcagtgggct	gcggcatacg	cctgcccac	600
ccagacactg	acctctactg	gttcaccctg	taccagtttt	tcctggcctt	tgccctgcct	660
tttgtggtca	tcacagccgc	atacgtgagg	atcctgcagc	gcatgacgtc	ctcagtggcc	720
ccgcctccc	agcgcagcat	ccggctgcgg	acaaagaggg	tgaccgcgac	agccatcgcc	780
atctgtctgg	tcttctttgt	gtgctgggca	ccctactatg	tgctacagct	gaccagttg	840
tccatcagcc	gcccagacct	cacctttgtc	tacttataca	atgcggccat	cagcttgggc	900
tatgccaaca	gctgcctcaa	cccctttgtg	tacatcgtgc	tctgtgagac	gttccgcaaa	960
cgcttggtcc	tgctcgtgaa	gcctgcagcc	caggggcagc	ttcgcgctgt	cagcaacgct	1020
cagacggctg	acgaggagag	gacagaaagc	aaaggcacct	gatac		1065

&lt;210&gt; 8

&lt;211&gt; 1704

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; MCH-1R antagonist binding protein

&lt;400&gt; 8

atggacctgg	aagcctcgtc	gctgcccact	ggtcccaatg	ccagcaacac	ctctgatggc	60
cccataacc	tcacttcggc	aggatcacct	cctcgcacgg	ggagcatctc	ctacatcaac	120
atcatcatgc	cttcggtggt	cggcaccatc	tgctcctcgtg	gcatcatcgg	gaactccacg	180
gtcatcttcg	cggtcgtgaa	gaagtccaag	ctgcactggg	gcaacaacgt	ccccgacatc	240
ttcatcatca	acctctcggg	agtagatctc	ctctttctcc	tgggcatgcc	cttcatgatc	300
caccagctca	tgggcaatgg	ggtgtggcac	tttggggaga	ccatgtgcac	cctcatcacg	360
gccatggatg	ccaatagtca	gttcaccagc	acctacatcc	tgaccgccat	ggccattgac	420
cgctacctgg	ccactgtcca	ccccatctct	tccacgaagt	tccggaagcc	ctctgtggcc	480
accttggtga	tctgcctcct	gtgggcccctc	tccttcatca	gcatcacccc	tgtgtggctg	540
tatgccagac	tcattcccctt	cccaggaggt	gcagtgggct	gcggcatacg	cctgcccac	600
ccagacactg	acctctactg	gttcaccctg	taccagtttt	tcctggcctt	tgccctgcct	660
tttgtggtca	tcacagccgc	atacgtgagg	atcctgcagc	gcatgacgtc	ctcagtggcc	720
ccgcctccc	agcgcagcat	ccggctgcgg	acaaagaggg	tgaccgcgac	agccatcgcc	780
atctgtctgg	tcttctttgt	gtgctgggca	ccctactatg	tgctacagct	gaccagttg	840
tccatcagcc	gcccagacct	cacctttgtc	tacttataca	atgcggccat	cagcttgggc	900
tatgccaaca	gctgcctcaa	cccctttgtg	tacatcgtgc	tctgtgaggt	cgacggtacc	960
gcggggccgg	gatccatcgc	caccatggtg	agcaaggggc	aggagctgtt	caccgggggtg	1020
gtgcccattc	tggtcgagct	ggacggcgac	gtaaaccggc	acaagtccag	cgtgtccggc	1080
gagggcgagg	gcgatgccac	ctacggcaag	ctgaccctga	agttcatctg	caccaccggc	1140
aagctgcccc	tgccctggcc	caccctcgtg	accaccctga	cctacggcgt	gcagtgtctc	1200
agccgctacc	ccgaccacat	gaagcagcac	gacttcttca	agtcgcgcat	gcccgaaggc	1260
tacgtccagg	agcgcaccat	cttcttcaag	gacgacggca	actacaagac	ccgcgccgag	1320
gtgaagtctg	agggcgacac	cctggtgaac	cgcactcagc	tgaaggccat	cgacttcaag	1380
gaggacggca	acatcctggg	gcacaagctg	gagtaacaact	acaacagcca	caacgtctat	1440
atcatggccg	acaagcagaa	gaacggcatc	aaggtgaact	tcaagatccg	ccacaacatc	1500
gaggacggca	gcgtgcagct	cgccgaccac	taccagcaga	acacccccat	cggcgacggc	1560
ccggtgctgc	tgcccgcaca	ccactacctg	agcaccagct	ccgccctgag	caaagacccc	1620
aacgagaagc	gcgatcacat	ggtcctgctg	gagttcgtga	ccgcgcgccc	gatcaactctc	1680
ggcatggacg	agctgtacaa	gtaa				1704

<210> 9  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer

<400> 9  
ccatggccat tgaccactac ctggccactg tcc 33

<210> 10  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer

<400> 10  
ggacagtggc caggtagtgg tcaatggcca tgg 33

<210> 11  
<211> 34  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer

<400> 11  
ctcttcacg aagttcgca agccctctgt ggcc 34

<210> 12  
<211> 34  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer

<400> 12  
ggccacagag ggcttcgca acttcgtgga agag 34

<210> 13  
<211> 47  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide primer

<400> 13  
tttgtgtaca tcgtgctctg tgaggtcgac ggtaccgcgg gcccggg 47

<210> 14  
 <211> 47  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Oligonucleotide primer

<400> 14  
 cccgggcccgc cggtagcggtc gacgtcacag agcagcatgt acacaaa

47

<210> 15  
 <211> 27  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Region between TM3 and TM4 in MCH-1R(i2/MC4R)

<400> 15  
 Asp Arg Tyr Phe Thr Ile Phe Tyr Ala Leu Gln Tyr His Asn Ile Met  
   1                  5                  10                  15  
 Thr Val Lys Arg Ala Thr Leu Val Ile Cys Leu  
                   20                  25

<210> 16  
 <211> 11  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> MCH-1R antagonist

<221> DISULFID  
 <222> (2)...(11)

<221> ACETYLTATION  
 <222> (1)...(1)

<221> AMIDATION  
 <222> (11)...(11)

<221> MOD\_RES  
 <222> (1)...(1)  
 <223> Xaa = des-amino-arginine

<221> MOD\_RES  
 <222> (9)...(10)  
 <223> Xaa = 5-aminovaleric acid

<400> 16  
 Xaa Cys Met Leu Gly Arg Val Tyr Xaa Xaa Cys  
   1                  5                  10



<210> 17  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> MCH-1R antagonist

<221> DISULFID  
<222> (2)...(11)

<221> ACETYLATION  
<222> (1)...(1)

<221> AMIDATION  
<222> (11)...(11)

<221> MOD\_RES  
<222> (1)...(1)  
<223> Xaa = des-amino-arginine

<221> MOD\_RES  
<222> (5)...(5)  
<223> Xaa = D-norleucine

<221> MOD\_RES  
<222> (9)...(10)  
<223> Xaa = 5-aminovaleric acid

<400> 17  
Xaa Cys Met Leu Xaa Arg Val Tyr Xaa Xaa Cys  
1 5 10

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**